

Datasheet for ABIN4886264  
**RICTOR ELISA Kit**



[Go to Product page](#)

## Overview

Quantity:	96 tests
Target:	RICTOR
Binding Specificity:	pThr1135, total
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA

## Product Details

Purpose:	Human Phospho-Rictor (T1135) and Total Rictor ELISA Kit. This assay semi-quantitatively measures phosphorylated Rictor (Thr1135) and Total Rictor in lysate samples.
Sample Type:	Cell Lysate, Tissue Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes Human Rictor phosphorylated at site Threonine-1135 and Total Rictor.
Characteristics:	<ul style="list-style-type: none"><li>• Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose)</li><li>• Screen numerous different cell lysates without performing a Western Blot analysis</li><li>• Minimal hands-on time, convenient, and non-radioactive material</li></ul>
Components:	<ul style="list-style-type: none"><li>• Pre-Coated 96-well Strip Microplate</li><li>• Wash Buffer</li></ul>

## Product Details

---

- Anti-Phospho Antibody
- Anti-Pan Antibody
- HRP-Conjugated Secondary Antibody
- Streptavidin-Conjugated HRP
- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

---

### Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2  $\mu$ L to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

## Target Details

---

Target: RICTOR

Alternative Name: Rictor ([RICTOR Products](#))

Gene ID: 253260

UniProt: [Q6R327](#)

Pathways: [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [Regulation of Actin Filament Polymerization](#), [CXCR4-mediated Signaling Events](#)

## Application Details

---

Sample Volume: 100  $\mu$ L

Plate: Pre-coated

Protocol:

1. Prepare all reagents and samples as instructed in the manual.
2. Add 100  $\mu$ L of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4  $^{\circ}$ C.
4. Add 100  $\mu$ L of prepared primary antibody to each well.
5. Incubate 1 h at RT.
6. Add 100  $\mu$ L of prepared 1X HRP-Streptavidin to each well.

## Application Details

---

7. Incubate 1 h at RT.
8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 µL of Stop Solution to each well.
11. Read at 450 nm immediately.

---

Restrictions: For Research Use only

## Handling

---

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

---

Expiry Date: 6 months