.-online.com antibodies

Datasheet for ABIN4886264 RICTOR ELISA Kit



Overview

Quantity:	96 tests
Target:	RICTOR
Binding Specificity:	pThr1135, total
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Human Phospho-Rictor (T1135) and Total Rictor ELISA Kit. This assay semi-quantitatively
	measures phophorylated Rictor (Thr1135) and Total Rictor in lysate samples.
Sample Type:	Cell Lysate, Tissue Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes Human Rictor phosphorylated at site
	Threonine-1135 and Total Rictor.
Characteristics:	 Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose)
	Screen numerous different cell lysates without performing a Western Blot analysis
	Minimal hands-on time, convenient, and non-radioactive material
Components:	Pre-Coated 96-well Strip Microplate
	Wash Buffer

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN4886264 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

	 Anti-Phospho Antibody Anti-Pan Antibody HRP-Conjugated Secondary Antibody Streptavidin-Conjugated HRP Assay Diluent TMB One-Step Substrate Stop Solution Lysis Buffer
	Positive Control Sample
Material not included:	Distilled or deionized water
	100 mL and 1 liter graduated cylinders
	Tubes to prepare sample dilutions
	Protease and Phosphatase inhibitors
	 Precision pipettes to deliver 2 µL to 1 mL volumes
	Adjustable 1-25 mL pipettes for reagent preparation
	Benchtop rocker or shaker
	Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	RICTOR
Alternative Name:	Rictor (RICTOR Products)
Gene ID:	253260
UniProt:	Q6R327
Pathways:	Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling
	Pathway, Regulation of Actin Filament Polymerization, CXCR4-mediated Signaling Events

Application Details

Sample Volume:	100 µL
Plate:	Pre-coated
Protocol:	 Prepare all reagents and samples as instructed in the manual. Add 100 μL of sample or positive control to each well. Incubate 2.5 h at RT or O/N at 4 °C. Add 100 μL of prepared primary antibody to each well. Incubate 1 h at RT. Add 100 μL of prepared 1X HRP-Streptavidin to each well.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN4886264 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

	 7. Incubate 1 h at RT. 8. Add 100 μL of TMB One-Step Substrate Reagent to each well. 9. Incubate 30 min at RT. 10. Add 50 μL of Stop Solution to each well. 11. Read at 450 nm immediately.
Restrictions:	For Research Use only
Handling	
Storage:	-20 °C
Storage Comment:	Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.
Expiry Date:	6 months