# antibodies -online.com





#### Go to Product pag

### Datasheet for ABIN4886285

#### SHP1 ELISA Kit

## Overview

Quantity:	96 tests
Target:	SHP1 (PTPN6)
Binding Specificity:	pSer591
Reactivity:	Human, Mouse
Method Type:	Sandwich ELISA
Application:	ELISA
Product Details	

Application:	ELISA
Product Details	
Purpose:	Human and Mouse Phospho-SHP1 (Ser591) ELISA Kit. This assay semi-quantitatively measures phophorylated SHP1 (Ser591) in lysate samples.
Sample Type:	Cell Culture Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human and Mouse SHP1 phosphorylated at site Serine-591.
Characteristics:	<ul> <li>Rapidly measure phosphorylated protein in lysates</li> <li>Screen numerous different cell lysates without performing a Western Blot analysis</li> <li>Minimal hands-on time, convenient, and non-radioactive material</li> </ul>
Components:	<ul> <li>Pre-Coated 96-well Strip Microplate</li> <li>Wash Buffer</li> <li>Anti-Phospho Antibody</li> <li>HRP-Conjugated Secondary Antibody</li> </ul>

#### **Product Details**

- · Assay Diluent
- · TMB One-Step Substrate
- · Stop Solution
- · Lysis Buffer
- · Positive Control Sample

#### Material not included:

- · Distilled or deionized water
- · 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- · Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- · Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

#### **Target Details**

Target:	SHP1 (PTPN6)
Alternative Name:	SHP1 (PTPN6 Products)
Gene ID:	5777
UniProt:	P29350, P29351
Pathways:	JAK-STAT Signaling, TCR Signaling, TLR Signaling, Nuclear Receptor Transcription Pathway, Positive Regulation of Peptide Hormone Secretion, Steroid Hormone Mediated Signaling Pathway, Response to Growth Hormone Stimulus, Regulation of Leukocyte Mediated Immunity, CXCR4-mediated Signaling Events, Signaling Events mediated by VEGFR1 and VEGFR2, BCR Signaling

#### **Application Details**

Sample Volume:	100 μL
Plate:	Pre-coated
Protocol:	Prepare all reagents and samples as instructed in the manual.
	2. Add 100 μL of sample or positive control to each well.
	3. Incubate 2.5 h at RT or O/N at 4 °C.
	4. Add 100 μL of prepared primary antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 μL of prepared 1X HRP-Streptavidin to each well.
	7. Incubate 1 h at RT.

#### **Application Details**

Expiry Date:

6 months

8. Add 100 µL of TMB One-Step Substrate Reagent to each well. 9. Incubate 30 min at RT. 10. Add 50 µL of Stop Solution to each well. 11. Read at 450 nm immediately. Restrictions: For Research Use only Handling Storage: -20 °C Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.