

Datasheet for ABIN4886291

## Phospholipase C gamma 2 ELISA Kit



[Go to Product page](#)

### Overview

Quantity:	96 tests
Target:	Phospholipase C gamma 2 (PLCG2)
Binding Specificity:	pThr277
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA

### Product Details

Purpose:	Human Phospho-SMAD4 (Thr277) ELISA Kit. This assay semi-quantitatively measures phosphorylated SMAD4 (Thr277) in lysate samples.
Sample Type:	Cell Lysate, Tissue Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human PLCG2 phosphorylated at site Threonine-277.
Characteristics:	<ul style="list-style-type: none"><li>• Rapidly measure phosphorylated protein in lysates</li><li>• Screen numerous different cell lysates without performing a Western Blot analysis</li><li>• Minimal hands-on time, convenient, and non-radioactive material</li></ul>
Components:	<ul style="list-style-type: none"><li>• Pre-Coated 96-well Strip Microplate</li><li>• Wash Buffer</li><li>• Anti-Phospho Antibody</li><li>• HRP-Conjugated Secondary Antibody</li></ul>

## Product Details

---

- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

---

### Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2  $\mu$ L to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

## Target Details

---

Target: Phospholipase C gamma 2 (PLCG2)

Alternative Name: PLCG2 ([PLCG2 Products](#))

Gene ID: 4089

UniProt: [Q13485](#)

Pathways: [RTK Signaling](#), [WNT Signaling](#), [Fc-epsilon Receptor Signaling Pathway](#), [Inositol Metabolic Process](#), [Myometrial Relaxation and Contraction](#), [Toll-Like Receptors Cascades](#), [VEGF Signaling](#), [BCR Signaling](#)

## Application Details

---

Sample Volume: 100  $\mu$ L

Plate: Pre-coated

Protocol:

1. Prepare all reagents and samples as instructed in the manual.
2. Add 100  $\mu$ L of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4  $^{\circ}$ C.
4. Add 100  $\mu$ L of prepared primary antibody to each well.
5. Incubate 1 h at RT.
6. Add 100  $\mu$ L of prepared 1X HRP-Streptavidin to each well.
7. Incubate 1 h at RT.
8. Add 100  $\mu$ L of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.

## Application Details

---

10. Add 50  $\mu$ L of Stop Solution to each well.
11. Read at 450 nm immediately.

---

Restrictions: For Research Use only

## Handling

---

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

---

Expiry Date: 6 months