# antibodies -online.com





# Datasheet for ABIN4886339

## **TYK2 ELISA Kit**

Go to Product page

	1//			

Quantity:	96 tests
Target:	TYK2
Binding Specificity:	pTyr1054, total
Reactivity:	Human, Mouse
Method Type:	Sandwich ELISA
Application:	ELISA
Product Details	
Purpose:	Human and Mouse Phospho-TYK2 (Tyr1054) and Total TYK2 ELISA Kit. This assay semi- quantitatively measures phophorylated TYK2 (Tyr1054) and Total TYK2 in lysate samples.
Sample Type:	Cell Culture Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human and Mouse TYK2 phosphorylated at site Tyrosine-1054 as well as total TYK2.
Characteristics:	<ul> <li>Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose)</li> <li>Screen numerous different cell lysates without performing a Western Blot analysis</li> <li>Minimal hands-on time, convenient, and non-radioactive material</li> </ul>
Components:	<ul><li>Pre-Coated 96-well Strip Microplate</li><li>Wash Buffer</li></ul>

### **Product Details**

- · Anti-Phospho Antibody
- · Anti-Pan Antibody
- HRP-Conjugated Secondary Antibody
- · Streptavidin-Conjugated HRP
- · Assay Diluent
- · TMB One-Step Substrate
- · Stop Solution
- · Lysis Buffer
- · Positive Control Sample

#### Material not included:

- Distilled or deionized water
- · 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- · Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- · Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

## **Target Details**

Target:	TYK2
Alternative Name:	TYK2 (TYK2 Products)
Gene ID:	7297
UniProt:	P29597, Q9R117
Pathways:	JAK-STAT Signaling, Hepatitis C

#### **Application Details**

Application Details		
Sample Volume:	100 μL	
Plate:	Pre-coated	
Protocol:	1. Prepare all reagents and samples as instructed in the manual.	
	2. Add 100 $\mu L$ of sample or positive control to each well.	
	3. Incubate 2.5 h at RT or O/N at 4 °C.	
	4. Add 100 μL of prepared primary antibody to each well.	
	5. Incubate 1 h at RT.	
	6. Add 100 μL of prepared 1X HRP-Streptavidin to each well.	
	7. Incubate 1 h at RT.	
	8. Add 100 µL of TMB One-Step Substrate Reagent to each well.	

## **Application Details**

	9. Incubate 30 min at RT.
	10. Add 50 μL of Stop Solution to each well.
	11. Read at 450 nm immediately.
Restrictions:	For Research Use only
Handling	
Storage:	-20 °C
Storage Comment:	Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of
	shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-
	Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell
	Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return
	unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20
	°C. Reconstituted Positive Control (Item K) should be stored at -70 °C.
Expiry Date:	6 months