

[Go to Product page](#)

## Datasheet for ABIN4886363

### PARP1 ELISA Kit

#### Overview

Quantity:	96 tests
Target:	PARP1
Binding Specificity:	Cleaved-Asp214, Cleaved-Gly215, total
Reactivity:	Human
Method Type:	Sandwich ELISA
Application:	ELISA

#### Product Details

Purpose:	Human Cleaved PARP (D214/G215) and PARP ELISA Kit. This ELISA is for measuring cleaved PARP (Asp-214/Gly-215) as well as PARP in human cell lysates.
Sample Type:	Cell Lysate, Tissue Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes human PARP cleaved at sites aspartic acid-214 and glycine-215.
Characteristics:	<ul style="list-style-type: none"><li>• Rapidly measures cleaved protein and pan protein in one experiment (for normalization purpose)</li><li>• Screen numerous different cell lysates without performing a Western Blot analysis</li><li>• Minimal hands-on time, convenient, and non-radioactive material</li></ul>
Components:	<ul style="list-style-type: none"><li>• Pre-Coated 96-well Strip Microplate</li><li>• Wash Buffer</li></ul>

## Product Details

- Anti-cleaved Antibody
- Anti-pan Antibody
- HRP-Conjugated Secondary Antibody
- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

### Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

## Target Details

Target: PARP1

Alternative Name: PARP1 ([PARP1 Products](#))

Gene ID: 142

UniProt: [P09874](#)

Pathways: [Apoptosis](#), [Caspase Cascade in Apoptosis](#), [DNA Damage Repair](#), [Production of Molecular Mediator of Immune Response](#), [Maintenance of Protein Location](#)

## Application Details

Sample Volume: 100 µL

Plate: Pre-coated

Protocol:

1. Prepare all reagents and samples as instructed in the manual.
2. Add 100 µL of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4 °C.
4. Add 100 µL of prepared primary antibody to each well.
5. Incubate 1 h at RT.
6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.
7. Incubate 1 h at RT.

## Application Details

---

8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 µL of Stop Solution to each well.
11. Read at 450 nm immediately.

---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

---

Storage:	-20 °C
----------	--------

Storage Comment:	Upon receipt, the kit should be stored at -20°C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4°C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge, and store at -20°C. Item D, store at 2-8°C for up to one month (store at -20°C for up to 6 months, avoid repeated freeze-thaw cycles). Reconstituted Positive Control (Item K) should be stored at -70°C.
------------------	---

---

Expiry Date:	6 months
--------------	----------