# antibodies -online.com







## anti-Nuclear Antigen antibody



#### Overview

Quantity:	50 μg
Target:	Nuclear Antigen (IPO-38)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Nuclear Antigen antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunoprecipitation (IP), ELISA, Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	Spleen cells of a human patient with hairy cell leukemia.
Clone:	Clon IPO-38
Specificity:	Reacts with human Proliferation Marker IPO-38. This antigen was detected after 12 h of PHA-induced activation in the early G1 phase but absent in non-stimulated lymphocytes.
Purification:	Purified (protein A)

#### **Target Details**

Target:	Nuclear Antigen (IPO-38)
Alternative Name:	Proliferation Marker IPO-38 (IPO-38 Products)

### **Application Details**

Application Notes:	Working dilution: Optimal dilutions should be determined by the end user.
	The following are guidelines only:
	ELISA (1:10000 - 1:50000) IHC - F, IHC - P (10 μg/mL) IP, WB (1:500 - 1:2000)
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.02 M potassium phosphate, 0.5 M NaCl, pH 7.2
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Long term: -20°C, Short term: -20°C