



[Go to Product page](#)

Datasheet for ABIN4889788

SHC1 ELISA Kit

3 Images

Overview

Quantity:	96 tests
Target:	SHC1
Binding Specificity:	pTyr427
Reactivity:	Human, Mouse, Rat
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Human, Mouse and Rat Phospho-SHC (Tyr427) ELISA Kit. This assay semi-quantitatively measures SHC phosphorylated at Tyrosine-427 in cell lysate samples.
Sample Type:	Cell Culture Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human, Mouse and Rat SHC phosphorylated at site Tyrosine-427.
Characteristics:	<ul style="list-style-type: none">• Rapidly measure phosphorylated protein in lysates• Screen numerous different cell lysates without performing a Western Blot analysis• Minimal hands-on time, convenient, and non-radioactive material
Components:	<ul style="list-style-type: none">• Pre-Coated 96-well Strip Microplate• Wash Buffer• Anti-Phospho Antibody• HRP-Conjugated Secondary Antibody

Product Details

- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target: SHC1

Alternative Name: SHC ([SHC1 Products](#))

Gene ID: 6464

UniProt: [P29353](#), [P98083](#), [O70143](#)

Pathways: [RTK Signaling](#), [TCR Signaling](#), [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [ER-Nucleus Signaling](#), [Signaling Events mediated by VEGFR1 and VEGFR2](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Sample Volume: 100 µL

Plate: Pre-coated

Protocol:

1. Prepare all reagents and samples as instructed in the manual.
2. Add 100 µL of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4 °C.
4. Add 100 µL of prepared primary antibody to each well.
5. Incubate 1 h at RT.
6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.
7. Incubate 1 h at RT.

Application Details

8. Add 100 μ L of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 μ L of Stop Solution to each well.
11. Read at 450 nm immediately.

Restrictions: For Research Use only


Handling

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.


Expiry Date: 6 months

Images




ELISA

Image 1. A431 cells were treated with EGF. Solubilize cells at 4×10^7 cells/ml in Cell Lysate Buffer. Serial dilutions of lysates were analyzed in this ELISA.



ELISA

Image 2. A431 cells were treated or untreated with EGF. Cell lysates were analyzed using this phosphoELISA and Western Blot.



ELISA

Image 3. A431 cells were treated or untreated with EGF. Cell lysates were analyzed using this phosphoELISA and Western Blot.