

Datasheet for ABIN4907330
anti-E-cadherin antibody



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	E-cadherin (CDH1)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This E-cadherin antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	This CADH1 antibody is generated from a mouse immunized with a recombinant protein between 1-392 amino acids from human CADH1.
Clone:	9C8
Isotype:	IgG1
Cross-Reactivity:	Human
Purification:	Purified by Protein G.

Target Details

Target:	E-cadherin (CDH1)
Alternative Name:	CDH1 (CDH1 Products)

Target Details



Background:	Synonyms: UVO, CDHE, ECAD, LCAM, Arc-1, CD324, Cadherin-1, CAM 120/80, Epithelial cadherin, E-cadherin, Uvomorulin, CDH1 Background: Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells, cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells. Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7. E-Cad/CTF2 promotes non-amyloidogenic degradation of Abeta precursors. Has a strong inhibitory effect on APP C99 and C83 production.
Gene ID:	999
UniProt:	P12830
Pathways:	WNT Signaling , Sensory Perception of Sound , Cell-Cell Junction Organization , Tube Formation

Application Details

Application Notes:	WB 1:300-5000 FCM 1:20-100 IHC-P 1:200-400 IF(IHC-P) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



Western Blotting

Image 1. Lane 1: A431 Cell lysates, Lane 2: T47D Cell lysates, probed with CADH1 (1579CT577.150.80) Monoclonal Antibody, unconjugated (bsm-51351M) at 1:4000 overnight at 4°C followed by a conjugated secondary antibody for 60 minutes at 37°C.

Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Paraformaldehyde-fixed, paraffin embedded human colon tissue, Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min, Block endogenous peroxidase by 3% hydrogen peroxide for 15 minutes, Blocking buffer (3% BSA) at room temperature for 30min, Antibody incubation with CADH1 (1579CT577.150.80) Monoclonal Antibody (bsm-51351M) at 1:25 for 1 hour at 37°C, followed by a conjugated secondary antibody for 20 minutes and DAB staining.