# .-online.com antibodies

Datasheet for ABIN4986886 GM-CSF ELISA Kit

Image



#### Overview

Quantity:	96 tests
Target:	GM-CSF (CSF2)
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	0.03-2.0 ng/mL
Minimum Detection Limit:	0.03 ng/mL
Application:	ELISA

## Product Details

Sample Type:	Cell Culture Supernatant, Serum, Plasma (heparin), Plasma (citrate), Plasma (EDTA)
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	Natural and recombinant Mouse GM-CSF Ligand
Sensitivity:	15 pg/mL
Material not included:	<ul> <li>Microplate reader.</li> <li>Pipettes and pipette tips.</li> <li>EP tube Deionized or distilled water.</li> </ul>

#### Target Details


Target:

GM-CSF (CSF2)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN4986886 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	GM-CSF (CSF2 Products)
Background:	Granulocyte macrophage colony-stimulating factor (GM-CSF), a 22 kDa glycosylated protein,
	was initially characterized as a factor that can support the in vitro colony formation of
	granulocytemacrophage progenitors. It is also a growth factor for erythroid, megakaryocyte,
	and eosinophil progenitors. GM-CSF is produced by a number of different cell types (including T
	cells, B cells, macrophages, mast cells, endothelial cells, fibroblasts, and adipocytes) in
	response to cytokine or inflammatory stimuli. On mature hematopoietic cells, GM-CSF is a
	survival factor for and activates the effector functions of granulocytes,
	monocytes/macrophages, and eosinophils (1, 2). GM-CSF promotes a Th1 biased immune
	response, angiogenesis, allergic inflammation, and the development of autoimmunity (3 5). It
	shows clinical effectiveness in ameliorating chemotherapyinduced neutropenia, and GM-CSF
	transfected tumor cells are utilized as cancer vaccines (6, 7). Mature mouse GM-CSF shares
	49 % -54 % amino acid sequence identity with canine, feline, human, and porcine GM-CSF and
	69 % with rat GM-CSF. GM-CSF exerts its biological effects through a heterodimeric receptor
	complex composed of GM-CSF Ra/CD116 and the signal transducing common $eta$ chain
	(CD131) which is also a component of the highaffinity receptors for IL-3 and IL-5 (8,9). In
	addition, GM-CSF binds a naturally occurring soluble form of GM-CSF Ra (10). The activity of
	GM-CSF is species specific between human and mouse. Mouse GM-CSF is only weakly active
	on rat cells, although rat GM-CSF is fully active on mouse cells (11).
Pathways:	JAK-STAT Signaling, Cellular Response to Molecule of Bacterial Origin

Application Notes:	Detection Wavelength: 450 nm
Sample Volume:	20 µL
Assay Time:	3 h
Plate:	Pre-coated
Restrictions:	For Research Use only
Londling	
Handling	
Storage:	4 °C



### ELISA

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN4986886 | 09/11/2023 | Copyright antibodies-online. All rights reserved.