

Datasheet for ABIN499239
anti-AIF antibody (C-Term)



[Go to Product page](#)

2 Images

Overview

Quantity:	0.1 mg
Target:	AIF (AIFM1)
Binding Specificity:	AA 593-606, C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AIF antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Enzyme Immunoassay (EIA)

Product Details

Immunogen:	Peptide corresponding to amino acids 593 to 606 of Human AIF .Remarks: This sequence is identical to those of Mouse and Rat AIF.
Isotype:	IgG
Specificity:	This antibody detects AIF at C-term.
Cross-Reactivity (Details):	Species reactivity (tested):Human, Mouse, Rat
Purification:	Affinity chromatography purified via peptide column

Target Details

Target:	AIF (AIFM1)
Alternative Name:	AIFM1 / AIF (AIFM1 Products)

Target Details

Background:	Apoptosis is characterized by several morphological nuclear changes including chromatin condensation and nuclear fragmentation. These changes are triggered by the activation of members of caspase family, caspase activated DNase, and several novel proteins (1). A novel gene, the product of which causes chromatin condensation and DNA fragmentation, was recently identified, cloned, and designated apoptosis inducing factor (AIF) (2). Like the critical molecules, cytochrome c and caspase-9, in apoptosis, AIF localizes in mitochondria. AIF translocates to the nucleus when apoptosis is induced and induces mitochondria to release the apoptogenic proteins cytochrome c and caspase-9. AIF induces chromatin condensation and DNA fragmentation, which are the hallmarks of apoptosis, of the isolated nucleus and the nucleus in live cells by microinjection. AIF is highly conserved between human and mouse and widely expressed (2).Synonyms: Apoptosis-inducing factor 1 mitochondrial, PDCD8, Programmed cell death protein 8
Gene ID:	9131
NCBI Accession:	NP_004199
UniProt:	O95831
Pathways:	Apoptosis , Positive Regulation of Endopeptidase Activity , Cell RedoxHomeostasis , Smooth Muscle Cell Migration , Warburg Effect

Application Details

Application Notes:	ELISA. Western blot: 0.25 to 1 µg/mL. A 67 kDa band should be detected. Immunocytochemistry. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------


Handling

Buffer:	PBS containing 0.02 % Sodium Azide as preservative
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C

Handling

Storage Comment: Store undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.

Images



Immunofluorescence

Image 1. Immunocytochemistry of AIF in K562 cells with AP30031PU-N AIF antibody at 5 µg/ml.

Western Blotting

Image 2. Western blot analysis of AIF in K562 with AP30031PU-N AIF antibody at (A) 0.5, (B) 1, and (C) 2 µg/ml.