.-online.com antibodies

Datasheet for ABIN4993358 Cyclin E1 ELISA Kit

Image



Overview

Quantity:	96 tests
Target:	Cyclin E1 (CCNE1)
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	78.125-5000 pg/mL
Minimum Detection Limit:	78.125 pg/mL
Application:	ELISA

Product Details

Purpose:	The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of CCNE1 in Mouse serum, plasma and other biological fluids
Sample Type:	Plasma, Serum
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This kit recognizes natural and recombinantMouseCCNE1. No significant cross-reactivity or interference between MouseCCNE1 and analogues was observed. Note: Limited by existing techniques, cross reaction may still exist, as it is impossible for us to complete the cross-reactivity detection between MouseCCNE1 and all the analogues.
Sensitivity:	46.875 pg/mL
Components:	Micro ELISA Plate Reference Standard

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN4993358 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

Product Details

	Concentrated Biotinylated Detection Ab
	Concentrated HRP Conjugate
	Reference Standard & Sample Diluent
	Biotinylated Detection Ab Diluent
	HRP Conjugate Diluent
	Concentrated Wash Buffer (25×)
	Substrate Reagent
	Stop Solutio
	Plate Sealer
	Product Description
	Certificate of Analysis
Material not included:	Microplate reader with 450nm wavelength filter
	High-precision transferpettor, EP tubes and disposable pipette tips
	37°C Incubator
	Deionized or distilled water
	Absorbent paper
	Loading slot for Wash Buffer

Target Details

Target:	Cyclin E1 (CCNE1)
Alternative Name:	Cyclin-E1 (CCNE1) (CCNE1 Products)
Pathways:	Cell Division Cycle, Intracellular Steroid Hormone Receptor Signaling Pathway, Nuclear
	Hormone Receptor Binding, Mitotic G1-G1/S Phases

Application Details


Comment:	Information on standard material:
	The formulation of the standard is 0.01 M PBS. The standard contains additives (1 $\%$ BSA).
	Information on reagents:
	Reagents include 1 M SO ₂ . Azide, thimerosal, 2-mercaptoethanol (2-ME) or any other poisonous
	materials are not used.
	Information on antibodies:
	The provided antibodies and their host vary in different kits. All antibodies are affinity purified

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN4993358 | 09/11/2023 | Copyright antibodies-online. All rights reserved. The sensitivity of this assay, or Lower Limit of Detection (LLD) was defined as the lowest protein concentration that could be differentiated from zero. It was determined by adding two standard deviations to the mean optical density value of twenty zero standard replicates and calculating the corresponding concentration.

Sample Volume:	100 μL
Plate:	Pre-coated
Protocol:	This ELISA kit uses Sandwich-ELISA as the method. The micro ELISA plate provided in this kit
	has been pre-coated with an antibody specific to CCNE1. Standards or samples are added to
	the appropriate micro ELISA plate wells and combined with the specific antibody. Then a
	biotinylated detection antibody specific for CCNE1and Avidin-Horseradish Peroxidase (HRP)
	conjugate is added to each micro plate well successively and incubated. Free components are
	washed away. The substrate solution is added to each well. Only those wells that contain
	CCNE1, biotinylated detection antibody and Avidin-HRP conjugate will appear blue in color. The
	enzyme-substrate reaction is terminated by the addition of a sulphuric acid solution and the
	color turns yellow. The optical density (OD) is measured spectrophotometrically at a
	wavelength of 450 nm \pm 2 nm. The OD value is proportional to the concentration of CCNE1.You
	can calculate the concentration of CCNE1in the samples by comparing the OD of the samples
	to the standard curve. 6th Edition, revised in June, 2015 www.elabscience.com 4
Restrictions:	For Research Use only
Handling	
Handling Advice:	All the reagents in the kit should be stored according to the labels on vials. Unused wells should
	be returned to the foil pouch with the desiccant pack and resealed along entire edge of zip-seal.
	Substrate Reagent shouldn't be kept at -20 °C (Check!). Exposure of reagents to strong light
	should be avoided in the process of incubation and storage. All the taps of reagents should be
	tightened to prevent evaporation and microbial contamination. If not to store reagents
	according to above suggestions, erroneous results may occur.
Storage:	4 °C/-20 °C
Storage Comment:	The unopened kit can be stored at 4°C for 1 month. If the kit is not used within 1 month,
	store the items separately according to the conditions since the kit is received.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN4993358 | 09/11/2023 | Copyright antibodies-online. All rights reserved.





ELISA

Image 1. Typical standard curve

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 4/4 | Product datasheet for ABIN4993358 | 09/11/2023 | Copyright antibodies-online. All rights reserved.