

## Datasheet for ABIN499541 anti-Caspase 1 antibody (C-Term)

# **Images**




_						
	1//	Д	rv	16	٦/	٨
U	W	$\vdash$	ΙV	Ιt	٦,	/V

Quantity:	0.1 mg
Target:	Caspase 1 (CASP1)
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Caspase 1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)
Product Details	
Product Details  Immunogen:	Human Casp-1 / ICE / IL-1b converting enzyme (C-Terminus) Peptide
	Human Casp-1 / ICE / IL-1b converting enzyme (C-Terminus) Peptide
Immunogen:	
Immunogen: Isotype:	IgG  Caspase-1 antibody was raised against a 16 amino acid peptide from near the carboxy-
Immunogen: Isotype: Specificity:	IgG  Caspase-1 antibody was raised against a 16 amino acid peptide from near the carboxy-terminus of human Caspase-1.
Immunogen: Isotype: Specificity: Purification:	IgG  Caspase-1 antibody was raised against a 16 amino acid peptide from near the carboxy-terminus of human Caspase-1.


## **Target Details**

9			
Background:	Caspases are a family of cysteine proteases that can be divided into the apoptotic and		
	inflammatory caspase subfamilies. Unlike the apoptotic caspases, members of the		
	inflammatory subfamily are generally not involved in cell death but are associated with the		
	immune response to microbial pathogens (reviewed in 1 and 2). Members of this subfamily		
	include caspase-1, -4, -5, and -12 and can activate proinflammatory cytokines such as IL-1? and		
	IL-18 (3,4). Caspase-1 was initially identified as an IL-1beta-converting enzyme (5), later		
	experiments revealed it to be a mammalian homolog of the C. elegans cell death gene ced-3 (6)		
	whose overexpression can induce apoptosis in fibroblasts (7). Synonyms: CASP-1, CASP1, ICE,		
	IL1BC, IL1BCE, Interleukin-1 beta convertase, Interleukin-1 beta-converting enzyme, p45		
Gene ID:	834		
NCBI Accession:	NP_150634		
UniProt:	P29466		
Pathways:	Apoptosis, Interferon-gamma Pathway, Positive Regulation of Endopeptidase Activity,		
	Inflammasome		
Application Details			
Application Notes:	ELISA. Western Blot: Casp-1 antibody can be used for the detection of Caspase-1 at 0.5 µg/mL.		
	Immunohistochemistry: Casp-1 antibody can be used for the detection of Caspase-1 at 0.5 $\mu$		
	g/mL.		
	Other applications not tested.		
	Optimal dilutions are dependent on conditions and should be determined by the user.		
Restrictions:	For Research Use only		
Handling			
Buffer:	PBS containing 0.02 % sodium azide.		
Preservative:	Sodium azide		
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		
	should be handled by trained staff only.		
Storage:	4 °C		
Storage Comment:	Store the antibody undiluted at 2-8 °C.		



### Immunohistochemistry (Paraffin-embedded Sections)

**Image 1.** Immunohistochemical staining of human heart tissue using AP30187PU-N Caspase-1 antibody at  $2 \mu g/ml$ .



### **Western Blotting**

**Image 2.** Western blot analysis of Caspase-1 in human heart cell lysate with AP30187PU-N Caspase-1 antibody at (A) 0.5, (B) 1, and (C)  $2 \mu g/ml$ .