



[Go to Product page](#)

Datasheet for ABIN4995886

anti-ALKBH1 antibody (AA 1-100) (Alexa Fluor 750)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | ALKBH1 |
| Binding Specificity: | AA 1-100 |
| Reactivity: | Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ALKBH1 antibody is conjugated to Alexa Fluor 750 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human ABH1/ALKB |
| Isotype: | IgG |
| Cross-Reactivity: | Mouse, Rat |
| Predicted Reactivity: | Human,Dog,Cow,Sheep,Pig,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | ALKBH1 |
| Alternative Name: | ABH1/ALKB (ALKBH1 Products) |

Target Details

Background: Synonyms: ABH, ABH1, alkB, AlkB, alkylation repair homolog 1 E. coli, ALKB1_HUMAN, ALKBH, ALKBH1, Alkylated DNA repair protein alkB homolog 1, Alkylation repair homolog 1, Alkylation repair, alkB homolog, Alpha-ketoglutarate-dependent dioxygenase ABH1, DNA lyase ABH1, hABH.

Background: ALKB protects DNA and RNA against damage from methylating compounds from the environment by directly reversing 1-methyladenine (1-meA) and 3-methylcytosine (3-meC) cytotoxic alkylation lesions in DNA and RNA. The enzymes act by oxidative demethylation, utilizing ferrous iron and alpha-ketoglutarate as cofactors, 2-oxoglutarate as a co-substrate, and molecular oxygen as the oxidizing agent. Deficiencies in DNA and RNA repair in mammals are associated with cancer, neurological disease and developmental defects. ALKB plays a role in resistance to anti-cancer drugs which attempt to damage tumor DNA. Escherichia coli ALKB protein belongs to the superfamily of 2-oxoglutarate- and iron(II)-dependent oxygenases.

Gene ID: 8846

Pathways: [DNA Damage Repair](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months