



[Go to Product page](#)

Datasheet for ABIN4996021

anti-CAPG antibody (AA 251-348) (AbBy Fluor® 680)

Overview

Quantity:	100 µL
Target:	CAPG
Binding Specificity:	AA 251-348
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CAPG antibody is conjugated to AbBy Fluor® 680
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Actin Regulatory Protein CAPG
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Horse
Purification:	Purified by Protein A.

Target Details

Target:	CAPG
Alternative Name:	Actin Regulatory Protein CAPG (CAPG Products)
Background:	Synonyms: Actin capping protein GCAP39, Actin regulatory protein CAP G, Actin regulatory

Target Details

protein CAP-G, AFCP, CAPG, CAPG_HUMAN, Capping protein actin filament gelsolin like, Capping protein gelsolin like, gCap39, Gelsolin like capping protein, Macrophage capping protein, Macrophage-capping protein, mbh1, MCP, Myc basic mot homolog 1.

Background: Caldesmon, Filamin 1, Nebulin, Plastin, ADF, Gelsolin, CapG, Dematin and Cofilin are differentially expressed Actin-binding proteins. Both muscular (CDh) and non-muscular (CD1) forms of Caldesmon bind to Actin as well as to Calmodulin and Myosin. CDh is expressed predominantly on thin filaments in smooth muscle, whereas CD1 is widely expressed in non-muscle tissues and cells. CapG, also designated Actin-regulatory protein and macrophage-capping protein, is a macrophage-specific protein that reversibly blocks the barbed ends of Actin filaments, but does not sever preformed ones. The interactions of CapG with Actin may be important in the regulation of nuclear and cytoplasmic structures. CapG is a calcium-sensitive DNA-binding protein that plays a role in macrophage function. It is expressed in macrophages and macrophage-like cells and can localize both to the nucleus and the cytoplasm.

Gene ID: 822

Pathways: [Regulation of Actin Filament Polymerization](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months