
antibodies - online.com







anti-AFAP1L2 antibody (AbBy Fluor® 750)



()	ve	K\ /		A .
	\cup	1 V/	-	V۷

Quantity:	100 μL
Target:	AFAP1L2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This AFAP1L2 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human AFAP1L2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat
Purification:	Purified by Protein A.

Target Details

Target:	AFAP1L2	
Alternative Name:	AFAP1L2 (AFAP1L2 Products)	
Background:	Synonyms: KIAA1914, XB130, Actin filament-associated protein 1-like 2, AF1L2_HUMAN,	
	AFAP1-like protein 2, afap1l2.	

Background: Actin filament associated protein (AFAP-110) interacts directly with actin filaments through its C-terminal actin-binding domain. AFAP-110 contains additional protein-binding domains as well, and serves as an adaptor protein. By linking signaling molecules to actin filaments, AFAP-110 provides a platform for the preparation of larger signaling complexes, activates Src kinases in response to cellular signals and also directly affects Actin organization as an Actin filament cross-linking protein. AFAP-1L2 (Actin filament-associated protein 1-like 2), also known as XB130, is a 818 amino acid cytoplasmic protein that contains two Pleckstrin homology (PH) domains, which are normally found in proteins involved in intracellular signaling. Like its relative AFAP110, AFAP-1L2 interacts with Src kinase and may play a role in Src-regulated transcription activation. AFAP-1L2 is expressed in thyroid and spleen and can also be detected at lower levels in lung, brain, pancreas and kidney. There are four isoforms of AFAP-1L2 that are produced as a result of alternative splicing events.

Pathways:

EGFR Signaling Pathway

Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	