antibodies .- online.com







anti-AKAP5 antibody (AA 611-714) (Alexa Fluor 750)



\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	AKAP5	
Binding Specificity:	AA 611-714	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This AKAP5 antibody is conjugated to Alexa Fluor 750	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human AKAP5	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Dog,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	AKAP5	
Alternative Name:	AKAP5 (AKAP5 Products)	

Target Details

Target Type:	Viral Protein		
Background:	Synonyms: A kinase PRKA anchor protein 5, A kinase anchor protein 5, A kinase anchor protein		
	79 kDa, A kinase anchoring protein 75/79, A-kinase anchor protein 5, A-kinase anchor protein 79		
	kDa, AKAP 5, AKAP 75, AKAP 79, AKAP-5, Akap5, AKAP5_HUMAN, AKAP75, AKAP79, cAMP		
	dependent protein kinase regulatory subunit II high affinity binding protein, cAMP-dependent		
	protein kinase regulatory subunit II high affinity-binding protein, H21.		
	Background: The type II cAMP-protein kinase (PKA) is a multifunctional kinase with a broad		
	range of substrates (1). Specificity of PKA signaling is thought to be mediated by the		
	compartmentalization of the kinase to specific sites within the cell. To maintain this specific		
	localization, the R subunit (RII) of PKA interacts with specific RII-anchoring proteins. This family		
	of proteins has been designated A-kinase anchoring proteins (AKAP) (1-3). Members of this		
	family, including MAP2 (microtubule-associated protein 2), neuronally expressed AKAP 79 and		
	AKAP 150, and the DNA binding AKAP 95, display differential tissue specificity and localization		
	(4-6). Evidence suggests that AKAP 79 and AKAP 150 are both capable of anchoring PKA to		
	postsynaptic densities (PSD), which are a network of proteins located on the internal surfaces		
	of excitatory synapses.		
Gene ID:	9495		
UniProt:	P24588		
Pathways:	cAMP Metabolic Process		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	