

Datasheet for ABIN4996411 anti-AMH antibody (AbBy Fluor® 750)



Go to Product page

| _ | | | | | |
|---|-----|---|----|-------------|-----|
| | 1// | r | Vİ | \triangle | ۸/ |
| | V | | VI | | / V |

| Quantity: | 100 μL |
|--------------|--|
| Target: | АМН |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This AMH antibody is conjugated to AbBy Fluor® 750 |
| Application: | Western Blotting (WB) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human AMH | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human | |
| Predicted Reactivity: | Mouse,Rat,Dog,Cow,Pig,Horse | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | AMH | |
|-------------------|---|--|
| Alternative Name: | AMH (AMH Products) | |
| Background: | Synonyms: MIS_HUMAN, Anti muellerian hormone, M, MIS, Muellerian inhibiting factor, | |
| | Mullerian inhibiting substance. | |

Target Details

Background: Anti mullerian hormone (AMH) is a member of the TGF beta superfamily. It is secreted as a homodimeric 140kD disulphide linked precursor that is cleaved to release the mature 30kD homodimer. Originally classified as a foetal testicular hormone that inhibits Mullerian duct development, AMH is expressed post natally by immature Sertoli cells, and to a lesser degree by granulosa cells. AMH plays a role in testicular differentiation and in the regulation of ovarian follicle growth.

Gene ID:

268

Pathways:

Negative Regulation of Hormone Secretion

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |