# antibodies .- online.com





CINED'S CONT.

#### Datasheet for ABIN4997063

## anti-Ataxin 2 antibody (AA 775-856) (Alexa Fluor 750)

Purified by Protein A.

Overview	
Quantity:	100 μL
Target:	Ataxin 2 (ATXN2)
Binding Specificity:	AA 775-856
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Ataxin 2 antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human ATX2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse,Chicken

## Target Details

Purification:

Target:	Ataxin 2 (ATXN2)
Alternative Name:	ATX2 (ATXN2 Products)

### **Target Details**

•			
Background:	Synonyms: ATX2, SCA2, ASL13, TNRC13, Ataxin-2, Spinocerebellar ataxia type 2 protein,		
	Trinucleotide repeat-containing gene 13 protein, ATXN2		
	Background: The autosomal dominant cerebellar ataxias (ADCA) are a heterogeneous group of		
	neurodegenerative disorders characterized by progressive degeneration of the cerebellum,		
	brain stem and spinal cord. Clinically, ADCA has been divided into three groups: ADCA types I-III.  ATX2 belongs to the autosomal dominant cerebellar ataxias type I (ADCA I) which are characterized by cerebellar ataxia in combination with additional clinical features like optic atrophy, ophthalmoplegia, bulbar and extrapyramidal signs, peripheral neuropathy and dementia. ATX2 is caused by expansion of a CAG repeat in the coding region of ATX2. Longer		
		expansions result in earlier onset of the disease. There are four named isoforms.	
		Gene ID:	6311
		UniProt:	Q99700
		Pathways:	Ribonucleoprotein Complex Subunit Organization
	Application Details		
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
Expiry Date:	12 months		