antibodies -online.com





Datasheet for ABIN4997350

anti-BEND2 antibody (Alexa Fluor 750)



()	1 /	0	rv	/ 1 /	71	Α.
	1//	-	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL	
Target:	BEND2	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This BEND2 antibody is conjugated to Alexa Fluor 750	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human BEND2/CXorf56
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

Target Details	
Target:	BEND2
Alternative Name:	BEND2 (BEND2 Products)
Background:	Synonyms: BEN domain-containing protein 2, BEN domain containing 2, BEND 2, BEND2, BEND-
	2, Chromosome X open reading frame 20, MGC33653, BEND2_HUMAN, CXorf56.
	Background: BEND2 is a 799 amino acid protein that contains two BEN domains. BEND2 exists

as two alternatively spliced isoforms and is considered a complete proteome. BEN domain

mediates proteinDNA and proteinprotein interactions during chromatin organization and transcription. BEN domain may play a role in organization of viral DNA during replication or transcription. The BEND2 gene maps to human chromosome Xp22.13. Chromosome X consists of about 153 million base pairs and nearly 1,000 genes. The X and Y chromosomes are the human sex chromosomes. The combination of an X and Y chromosome lead to normal male development while two copies of X lead to normal female development. Color blindness, hemophilia, and Duchenne muscular dystrophy are well known X chromosome-linked conditions which affect males more frequently as males carry a single X chromosome.

Application Details

Storage:

Expiry Date:

Storage Comment:

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

-20 °C

12 months