antibodies .- online.com







anti-COPB1 antibody (AA 531-630) (Alexa Fluor 750)



\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	COPB1	
Binding Specificity:	AA 531-630	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This COPB1 antibody is conjugated to Alexa Fluor 750	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human beta COP	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Chicken	
Purification:	Purified by Protein A.	

Target Details

Target:	COPB1
Alternative Name:	beta COP (COPB1 Products)

Target Details

Bac	kar	'nΙ	ın	Ч.
Duo		\sim	<i>.</i>	ч.

Synonyms: Beta-coat protein, betacop, Coatomer beta subunit, Coatomer protein complex subunit beta 1, Coatomer protein complex subunit beta, Coatomer subunit beta, COPB, COPB1, DKFZp761K102, FLJ10341.

Background: The Golgi complex is a key organelle where processing and sorting of newly synthesized proteins occurs. Membrane traffic from the endoplasmic reticulum (ER) to the Golgi complex and from the Golgi complex to the different final cellular destinations is believed to be mediated by carrier vesicles. Two populations of coated vesicles mediate biosynthetic membrane traffic between the different membrane-bound compartments. Clathrin-coated vesicles carry proteins to endocytic organelles and secretory granules, whilst non-clathrin-coated vesicles are involved in intra-Golgi transport and transport from the ER to the Golgi complex. Beta COP is a member of a set of protein which are believed to associate with the non-clathrin coated vesicles. Golgi-derived non-clathrin-coated vesicles are believed to act as bulk carriers, whereas clathrin-coated vesicles carry a selective cargo of membrane proteins.

Gene ID:

1315

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months