

Datasheet for ABIN4997465

anti-Biglycan antibody (AA 101-200) (AbBy Fluor® 750)



Overview

Alternative Name:

Overview	
Quantity:	100 μL
Target:	Biglycan (BGN)
Binding Specificity:	AA 101-200
Reactivity:	Human, Rat, Mouse, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Biglycan antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human Biglycan
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rabbit, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.
Target Details	
Target:	Biglycan (BGN)
Altawaati ya Nlawaa	Disk or as (DON Day don't)

Biglycan (BGN Products)

Target Details

•	
Background:	Synonyms: PGI, DSPG1, PG-S1, SLRR1A, Biglycan, Bone/cartilage proteoglycan I, BGN
	Background: The protein encoded by this gene is a small cellular or pericellular matrix
	proteoglycan that is closely related in structure to two other small proteoglycans, decorin and
	fibromodulin. The encoded protein and decorin are thought to be the result of a gene
	duplication. Decorin contains one attached glycosaminoglycan chain, while this protein
	probably contains two chains. For this reason, this protein is called biglycan. This protein plays
	a role in assembly of collagen fibrils and muscle regeneration. It interacts with several proteins
	involved in muscular dystrophy, including alpha-dystroglycan, alpha- and gamma-sarcoglycan
	and collagen VI, and it is critical for the assembly of the dystrophin-associated protein complex.
Gene ID:	633
UniProt:	P21810
Pathways:	Glycosaminoglycan Metabolic Process
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months