

Datasheet for ABIN4997481

anti-KCNMA1 antibody (AA 21-120) (AbBy Fluor® 750)



Go to Product page

V-	r\/	ID	۱۸/

Overview		
Quantity:	100 μL	
Target:	KCNMA1	
Binding Specificity:	AA 21-120	
Reactivity:	Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This KCNMA1 antibody is conjugated to AbBy Fluor® 750	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human BKCA alpha	
Isotype:	lgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Cow,Pig,Chicken,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	KCNMA1	

Target Details

Background:	Synonyms: subfamily M subunit alpha-1, BK channel, BKCA alpha, BKCA alpha subunit, BKTM,	
	Calcium-activated potassium channel, Calcium-activated potassium channel subunit alpha-1,	
	Drosophila slowpoke like, hSlo, KVCAalpha, KCa1.1, KCMA1_HUMAN, KCNMA, KCNMA1, Maxi	
	K channel, Maxi Potassium channel alpha, MaxiK, SAKCA, SLO alpha, SLO, Slo homolog, Slo-	
	alpha, Slo1, Slowpoke homolog.	
	Background: MaxiK channels are large conductance, voltage and calcium-sensitive potassium	
	channels which are fundamental to the control of smooth muscle tone and neuronal	
	excitability. MaxiK channels can be formed by 2 subunits: the pore-forming alpha subunit, which	
	is the product of this gene, and the modulatory beta subunit. Intracellular calcium regulates the	
	physical association between the alpha and beta subunits. Alternatively spliced transcript	
	variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008].	
Pathways:	Regulation of Hormone Metabolic Process, Sensory Perception of Sound	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	