# antibodies - online.com







# anti-BOP1 antibody (AA 201-300) (Alexa Fluor 680)



( )	1 /	$\sim$	KI /	110	Νę
	1//	$\vdash$	I \/	1 ←	٠// ٢

Quantity:	100 μL	
Target:	BOP1	
Binding Specificity:	AA 201-300	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This BOP1 antibody is conjugated to Alexa Fluor 680	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human BOP1	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Chicken	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	BOP1
Alternative Name:	BOP1 (BOP1 Products)

## **Target Details**

Expiry Date:

rarget Details		
Background:	Synonyms: AU020183, AW146150, Block of proliferation 1, Block of proliferation 1 protein,	
	Bop1, Bop1 block of proliferation 1, BOP1_HUMAN, D18861, Erb1p, Kiaa0124, MGC109114,	
	mKIAA0124, Ribosome biogenesis protein BOP1.	
	Background: Predominantly localized to the nucleolus, BOP1 (Block of proliferation 1 protein) is	
	a 746 amino acid highly conserved non-ribosomal protein that is involved in ribosome	
	biogenesis. Truncation of the amino terminus of BOP1 leads to cell growth arrest in the G1	
	phase and specific inhibition of 28S and 5.8S rRNA synthesis, as well as a deficit in the	
	cytosolic 60S ribosomal subunit. This suggests that BOP1 is involved in the formation of	
	mature rRNAs and in the biogenesis of the 60S ribosomal subunit. BOP1 physically interacts	
	with pescadillo (a protein involved in cell proliferation) and enables efficient incorporation of	
	pescadillo into the nucleolar preribosomal complexes, thereby affecting rRNA maturation and	
	the cell cycle. The BOP1-pescadillo complex is also necessary for biogenesis of 60S ribosomal	
	subunits. Deregulation of BOP1 may lead to colorectal tumorigenesis.	
Gene ID:	23246	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Francisco Deter	10	

12 months