antibodies -online.com





anti-BTBD17 antibody (AA 101-200) (Alexa Fluor 680)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	BTBD17	
Binding Specificity:	AA 101-200	
Reactivity:	Mouse, Human, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This BTBD17 antibody is conjugated to Alexa Fluor 680	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human BTBD17	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Sheep	
Purification:	Purified by Protein A.	

Target Details

Target:	BTBD17
Alternative Name:	BTBD17 (BTBD17 Products)

Target Details

Background:	Synonyms: BTB/POZ domain-containing protein 17, btbd17, BTBDH_HUMAN, Galectin-3-			
	binding protein-like.			
	Background: The BTB (broad-complex, Tramtrack and Bric a brac) domain, also known as the			
	POZ (Poxvirus and zinc finger) domain, is an N-terminal homodimerization domain that			
	contains multiple copies of kelch repeats and/or C2H2-type zinc fingers. Proteins that contain			
	BTB domains are thought to be involved in transcriptional regulation via control of chromatin			
	structure and function. BTBD17 (BTB/POZ domain-containing protein 17), also known as			
	BTBD17A, galectin-3-binding protein-like or LGALS3BPL, is a 478 amino acid protein that			
	contains one BTB (POZ) domain and a BACK (BTB/Kelch associated) domain. The gene			
	encoding BTBD17 maps to human chromosome 17, which comprises over 2.5 % of the human			
	genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with			
	chromosome 17, namely, p53 and BRCA1. Malfunction or loss of p53 expression is associated			
	with malignant cell growth and Li-Fraumeni syndrome.			
Gene ID:	388419			
UniProt:	A6NE02			
Application Details				
Application Notes:	IF(IHC-P) 1:50-200			
	IF(IHC-F) 1:50-200			
	IF(ICC) 1:50-200			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	1 μg/μL			
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and			
	50 % Glycerol.			
Preservative:	ProClin			
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be			
	handled by trained staff only.			
Storage:	-20 °C			
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.			

	lI	1:
-	เวทก	111111
	land	11110

Expiry Date:

12 months