antibodies -online.com





anti-AAMDC antibody (Alexa Fluor 680)



	_	100		
Go to	n Pro	אלוומ	nt na	ane

_							
0	V	е	r١	/1	е	V	1

Overview		
Quantity:	100 μL	
Target:	AAMDC	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This AAMDC antibody is conjugated to Alexa Fluor 680	
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human C110RF67/PTD015	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Purified by Protein A.	
Target Details		
Target:	AAMDC	
Alternative Name:	C110RF67 (AAMDC Products)	
Background:	Synonyms: Chromosome 11 open reading frame 67, CK067, FLJ21035, Hypothetical protein LOC28971, MGC3367, PTD015, UPF0366 protein C11orf67, AAMDC_HUMAN. Background: PTD015 is a 122 amino acid protein that belongs to the UPF0366 family. Existing	

as three alternatively spliced isoforms, the PTD015 gene is conserved in dog, cow, mouse, rat,

chicken and zebrafish, and maps to human chromosome 11q14.1. With approximately 135 million base pairs and 1,400 genes, chromosome 11 makes up around 4 % of human genomic DNA and is considered a gene and disease association dense chromosome. The chromosome 11 encoded Atm gene is important for regulation of cell cycle arrest and apoptosis following double strand DNA breaks. Atm mutation leads to the disorder known as ataxia-telangiectasia. The blood disorders Sickle cell anemia and thalassemia are caused by HBB gene mutations. Wilms' tumors, WAGR syndrome and Denys-Drash syndrome are associated with mutations of the WT1 gene. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are also associated with defects in chromosome 11. The PTD015 gene product has been provisionally designated PTD015 pending further characterization.

UniProt:

Q9H7C9

IF(IHC-P) 1:50-200

Application Details

Application Notes:

Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months