

## Datasheet for ABIN4999034

## anti-NLRP1 antibody (AA 301-400) (AbBy Fluor® 680)



Overview	
Quantity:	100 μL
Target:	NLRP1
Binding Specificity:	AA 301-400
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NLRP1 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human CARD7/NALP1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat
Purification:	Purified by Protein A.
Target Details	
Target:	NLRP1
Alternative Name:	CARD7 (NLRP1 Products)

Background:

Synonyms: LRR and PYD domains-containing protein 1, CARD 7, CARD7, Caspase recruitment domain protein 7, Caspase recruitment domain-containing protein 7, CLR17.1, Death effector filament forming Ced 4 like apoptosis protein, Death effector filament-forming ced-4-like apoptosis protein, DEFCAP, DEFCAP L/S, NAC alpha/beta/gamma/delta, NAC, NACHT, NACHT leucine rich repeat and PYD containing 1, NACHT leucine rich repeat and PYD pyrin domain containing 1, NACHT leucine rich repeat and pyrin domain containing 1, NACHT LRR and PYD containing protein 1, NALP 1, NALP1, NALP1\_HUMAN, NLR family pyrin domain containing 1, NLRP 1, NLRP1, NLRP1 protein, Nucleotide binding domain and caspase recruitment domain, Nucleotide binding oligomerization domain leucine rich repeat and pyrin domain containing 1, Nucleotide-binding domain and caspase recruitment domain, PP 1044, PP1044. Background: This gene encodes a member of the Ced-4 family of apoptosis proteins. Cedfamily members contain a caspase recruitment domain (CARD) and are known to be key mediators of programmed cell death. The encoded protein contains a distinct N-terminal pyrinlike motif, which is possibly involved in protein-protein interactions. This protein interacts strongly with caspase 2 and weakly with caspase 9. Overexpression of this gene was demonstrated to induce apoptosis in cells. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene, but the biological validity of some variants has not been determined. [provided by RefSeq, Jul 2008].

Gene ID: 22861

Pathways: Positive Regulation of Endopeptidase Activity, Inflammasome

**Application Details** 

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration:  $1 \mu g/\mu L$ 

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and

50 % Glycerol.

Preservative: ProClin

## Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months