



[Go to Product page](#)

Datasheet for ABIN4999072

anti-CASK antibody (AA 651-750) (Alexa Fluor 680)

Overview

Quantity:	100 µL
Target:	CASK
Binding Specificity:	AA 651-750
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CASK antibody is conjugated to Alexa Fluor 680
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CASK
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CASK
Alternative Name:	CASK (CASK Products)

Target Details

Background: Synonyms: CAGH39, Caki, Calcium/calmodulin dependent serine protein kinase, Calcium/calmodulin dependent serine protein kinase MAGUK family, Calcium/calmodulin dependent serine protein kinase membrane associated guanylate kinase, Calcium/calmodulin-dependent serine protein kinase, CAMGUK, CAMGUK protein, CAMGUK, drosophila, homolog of antibody casK, CMG, CSKP_HUMAN, DXPri1, DXRib1, FGS4, hCASK, LIN 2, Lin 2 homolog, LIN2 antibody Lin2 homolog, MICPCH, MRXSNA, Pals3, Peripheral plasma membrane protein CASK, Protein lin-2 homolog, TNRC8, Trinucleotide repeat containing 8, Vertebrate LIN2 homolog.

Background: The MAGUK (membrane-associated guanylate kinase homologs) family of proteins contain multiple protein-binding domains and are involved in cell junction organization, tumor suppression, and signaling. CASK (also designated LIN-2) belongs to a MAGUK subfamily which is characterized by a novel domain structure that consists of a calcium/calmodulin-dependent protein kinase domain followed by PDZ, SH3 and guanylate kinase-like (GUK) domains. CASK is expressed in rat brain where it binds to cell-surface proteins, such as neurexin and syndecan, and is thought to be involved in signaling at neuronal synapses. CASK translocates to the nucleus and interacts with Tbr-1 to form a complex, which binds to a specific DNA sequence (the T-element), and induces the expression of specific genes, including Reelin. CASK displays a transcription regulation function, which appears crucial for cerebrocortical development.

Pathways: [Synaptic Vesicle Exocytosis](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months