

Datasheet for ABIN4999155

anti-Caveolin 2 antibody (pTyr19) (AbBy Fluor® 750)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Caveolin 2 (CAV2)
Binding Specificity:	pTyr19
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Caveolin 2 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from mouse Caveolin2 around the phosphorylation site of Tyr19.
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	Caveolin 2 (CAV2)
Alternative Name:	Caveolin-2 (CAV2 Products)

Target Details

Background:	Synonyms: Caveolin 2 phospho Y19, p-Caveolin 2phospho Y19, CAV, CAV2, CAV2_HUMAN, Caveolae protein 20 kD, Caveolin 2, Caveolin 2 isoform a and b, Caveolin 2 isoform c, Caveolin-2, MGC12294, CAV2_HUMAN. Background: Cellular localization: Nucleus. Cytoplasm. Golgi apparatus membrane. Cell membrane. Membrane > caveola. Potential hairpin-like structure in the membrane. Membrane protein of caveolae. Tyr-19-phosphorylated form is enriched at sites of cell-cell contact and is translocated to the nucleus in complex with MAPK1 in response to insulin (By similarity). Tyr-27-phosphorylated form is located both in the cytoplasm and plasma membrane. CAV1-mediated Ser-23-phosphorylated form locates to the plasma membrane. Ser-36-phosphorylated form resides in intracellular compartments.
Gene ID:	858
Pathways:	Regulation of G-Protein Coupled Receptor Protein Signaling , Skeletal Muscle Fiber Development

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months