## antibodies -online.com





## anti-GRIK1 antibody (C-Term)

2 Images



Go to Product page

$\sim$	
( )\/⊝	view
$\bigcirc$ $\lor$ $\bigcirc$	V I C V V

Background:


Quantity:	0.1 mg
Target:	GRIK1
Binding Specificity:	C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)
Product Details	
lmmunogen:	Grik1 antibody was raised against a 15 amino acid peptide near the carboxy terminus of the human Grik1.
Isotype:	IgG
Purification:	Affinity chromatography purified via peptide column
Target Details	
Target:	GRIK1
Alternative Name:	Glutamate Receptor 5 / GLUR5 (GRIK1 Products)

Glutamate receptors are the predominant excitatory neurotransmitter receptors in the

mammalian brain and are activated in a variety of normal neurophysiologic processes. Grik1,


also known as glutamate receptor 5, belongs to the kainate family of glutamate receptors,

	which are composed of four subunits and function as ligand-activated ion channels. Grik1 is
	expressed in GABAergic interneurons of the hippocampus and are thought to participate in the
	formation of various subtypes of kainate receptors with Grik2 and KA2. Stimulation of Grik1
	leads to intracellular calcium release and activation of protein kinase C. Excessive activation
	has been associated with psychiatric, neurological and neurodegenerative diseases. Numerous
	isoforms of Grik1 are known to exist and may be subject to RNA editing within the second
	transmembrane domain, which is thought to alter the properties of ion flow. Synonyms: EAA3,
	Excitatory amino acid receptor 3, GRIK1, Glutamate receptor ionotropic kainate 1
Gene ID:	2897
UniProt:	P39086
Pathways:	Synaptic Membrane, Regulation of long-term Neuronal Synaptic Plasticity
Application Details	
Application Notes:	ELISA. Western Blot: 0.5 - 1 μg/mL. Immunohistochemistry.
	Other applications not tested.
	Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only
Handling	
Buffer:	PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C.



## **Immunohistochemistry (Paraffin-embedded Sections)**

**Image 1.** Immunohistochemical staining of human brain tissue using AP30380PU-N Grik1 antibody at  $2.5 \, \mu g/ml$ .



## **Western Blotting**

Image 2. Western blot analysis of Grik1 in P815 cell lysate with AP30380PU-N Grik1 antibody at (A) 0.5, (B) 1 and (C) 2  $\mu$ g/ml.