antibodies -online.com





anti-CCDC138 antibody (AA 551-665) (Alexa Fluor 750)



Overview	
Overview	

Quantity:	100 μL
Target:	CCDC138
Binding Specificity:	AA 551-665
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCDC138 antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCDC138
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CCDC138
Alternative Name:	CCDC138 (CCDC138 Products)

Target Details

Background:	Synonyms: CCDC 138, CCDC138, Coiled coil domain containing 138, Coiled coil domain		
	containing protein 138, coiled-coil domain containing 138, FLJ 32745, FLJ32745,		
	CC138_HUMAN.		
	Background: CCDC138, also known as FLJ32745, is a 685 amino acid protein expressed as two		
	isoforms produced by alternative splicing. The gene that encodes CCDC138 maps to human		
	chromosome 2q12.3. The second largest human chromosome, chromosome 2 consists of 237		
	million bases, encodes over 1,400 genes and makes up approximately 8 % of the human genome. A number of genetic diseases are linked to genes on chromosome 2. Harlequin icthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene. The lipid metabolic disorder sitosterolemia is associated with ABCG5 and ABCG8. An extremely rare recessive genetic disorder, Alstr syndrome is due to mutations in the ALMS1 gene. Interestingly, chromosome 2 contains what appears to be a vestigial second centromere and		
			vestigial telomeres which gives credence to the hypothesis that human chromosome 2 is the
			result of an ancient fusion of two ancestral chromosomes seen in modern form today in apes.
Gene ID:		165055	
Pathways:	BCR Signaling		
Application Details			
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months