

Datasheet for ABIN4999270

**anti-CCDC50 antibody (AA 251-306) (AbBy Fluor® 680)**[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	CCDC50
Binding Specificity:	AA 251-306
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCDC50 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCDC50
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat
Purification:	Purified by Protein A.

## Target Details

Target:	CCDC50
Alternative Name:	CCDC50 ( <a href="#">CCDC50 Products</a> )

## Target Details

---

**Background:** Synonyms: C3orf6, C3orf6 YMER, CCD50\_HUMAN, CCDC 50, Ccdc50, Chromosome 3 open reading frame 6, Coiled coil domain containing 50, Coiled coil domain containing protein 50, Coiled-coil domain-containing protein 50, Protein Ymer, Ymer, Ymer protein.

Background: This gene encodes a soluble, cytoplasmic, tyrosine-phosphorylated protein with multiple ubiquitin-interacting domains. Mutations in this gene cause nonsyndromic, postlingual, progressive sensorineural DFNA44 hearing loss. In mouse, the protein is expressed in the inner ear during development and postnatal maturation and associates with microtubule-based structures. This protein may also function as a negative regulator of NF- $\kappa$ B signaling and as an effector of epidermal growth factor (EGF)-mediated cell signaling. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Oct 2008].

---

**Gene ID:** 152137

## Application Details

---

**Application Notes:** IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

---

**Concentration:** 1  $\mu$ g/ $\mu$ L

---

**Buffer:** Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

**Preservative:** ProClin

---

**Precaution of Use:** This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

**Storage:** -20 °C

---

**Storage Comment:** Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

**Expiry Date:** 12 months