# antibodies -online.com





## anti-CCDC54 antibody (AA 221-328) (Alexa Fluor 680)



Go to Product page

$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	CCDC54	
Binding Specificity:	AA 221-328	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CCDC54 antibody is conjugated to Alexa Fluor 680	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCDC54	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	CCDC54
Alternative Name:	CCDC54 (CCDC54 Products)

Precaution of Use:

Storage Comment:

Storage:

Target Details	
Background:	Synonyms: Coiled-coil domain containing 54, FLJ25362, NYD-SP17, Testes development-
	related NYD-SP17, Testis development protein NYD-SP17.
	Background: The specific function of CCDC54 (coiled-coil domain containing 54) is not yet
	known. The CCDC54 gene appears to be conserved in chimpanzee, dog, mouse, and
	rat.CCDC54, also known as coiled-coil domain-containing protein 54 or testis development
	protein NYD-SP17, is a 328 amino acid protein that is phosphorylated on threonine 182 during
	post-translational modification. The gene encoding NYD-SP17 maps to human chromosome
	3q13.12 and mouse chromosome 16 B5. Human chromosome 3 houses over 1,100 genes,
	including a chemokine receptor (CKR) gene cluster and a variety of human cancer-related gene
	loci. Key tumor suppressing genes on chromosome 3 include those that encode the apoptosis
	mediator RASSF1, the cell migration regulator HYAL1 and the angiogenesis suppressor
	SEMA3B. Marfan Syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta
	and Charcot-Marie-Tooth disease are a few of the numerous genetic diseases associated with
	chromosome 3.
Gene ID:	84692
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
D (11	The state of the police of the

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

handled by trained staff only.

-20 °C

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

$\vdash$	land	lına
	iaria	шц

Expiry Date:

12 months