antibodies - online.com







anti-CCDC98 antibody (AA 51-150) (Alexa Fluor 680)



| () | 11/ | IN | /ie | A . |
|-----|-------------------|------|--------|--------|
| | / // | ۱ ات | / (− | ' \/\/ |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | CCDC98 (FAM175A) | |
| Binding Specificity: | AA 51-150 | |
| Reactivity: | Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This CCDC98 antibody is conjugated to Alexa Fluor 680 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human CCDC98 | |
|-----------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Mouse | |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | CCDC98 (FAM175A) |
|-------------------|---------------------------|
| Alternative Name: | CCDC98 (FAM175A Products) |

Target Details

| Background: | Synonyms: ABRA1, CCDC98, BRCA1-A complex subunit Abraxas, Coiled-coil domain-containing | | |
|---------------------|---|--|--|
| | protein 98, Protein FAM175A, FAM175A, UNQ496/PRO1013 | | |
| | Background: Involved in DNA damage response and double-strand break (DSB) repair. | | |
| | Component of the BRCA1-A complex, acting as a central scaffold protein that assembles the | | |
| | various components of the complex and mediates the recruitment of BRCA1. The BRCA1-A | | |
| | complex specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA | | |
| | lesion sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at DSBs. | | |
| | This complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked | | |
| | ubiquitin on histones H2A and H2AX. | | |
| Gene ID: | 84142 | | |
| UniProt: | Q6UWZ7 | | |
| Pathways: | DNA Damage Repair, Positive Regulation of Response to DNA Damage Stimulus | | |
| Application Details | | | |
| Application Notes: | IF(IHC-P) 1:50-200 | | |
| | IF(IHC-F) 1:50-200 | | |
| | IF(ICC) 1:50-200 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | | |
| | 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | | |
| | handled by trained staff only. | | |
| Storage: | -20 °C | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | |
| Expiry Date: | 12 months | | |
| | | | |