

Datasheet for ABIN499964 anti-IKAP/p150 antibody (C-Term)

2 Images



Go to Product page

_				
()	ve.	rv/	101	Λ

Quantity:	0.1 mg	
Target:	IKAP/p150 (ELP1)	
Binding Specificity:	C-Term	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This IKAP/p150 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunofluorescence (IF), Enzyme Immunoassay (EIA)	
Product Details		
Immunogen:	IKAP antibody was raised against a 16 amino acid peptide from near the carboxy terminus of human IKAP.	
anogen.		
Isotype:		
	human IKAP.	
Isotype:	human IKAP.	
Isotype: Specificity:	human IKAP. IgG This antibody reacts to IKAP.	
Isotype: Specificity: Purification:	human IKAP. IgG This antibody reacts to IKAP.	
Isotype: Specificity: Purification: Target Details	human IKAP. IgG This antibody reacts to IKAP. Affinity chromatography purified via peptide column	
Isotype: Specificity: Purification: Target Details Target:	human IKAP. IgG This antibody reacts to IKAP. Affinity chromatography purified via peptide column IKAP/p150 (ELP1)	

to IKKalpha, IKKbeta, NF-kappaB, and the NF-kappaB-inducing kinase (NIK), although later evidence has cast doubt on this. More recent reports show that mutations in IKAP such as a frameshift leading to a truncated protein or a missense mutation that leads to defective phosphorylation are responsible for the autosomal recessive genetic disease familial dysautonomia (FD). Reports indicating that it forms part of the RNA polymerase II transcription elongation complex suggest that this disease may be due to compromised transcription elongation. More recently, it was shown that IKAP associates with c-Jun N-terminal kinase (JNK) and could specifically enhance JNK activation induced by the upstream JNK activators MEKK1 and ASK1, indicating another possible cause for FD. At least two isoforms of IKAP are known two exist. Synonyms: Elongator complex protein 1, IKAP, IKK complex-associated protein, IkappaB kinase complex-associated protein, p150

Gene ID:

8518

UniProt:

095163

Application Details

Application Notes:

ELISA. Western Blot: 0.5 to 1 µg/mL. Immunocytochemistry.

Other applications not tested.


Optimal dilutions are dependent on conditions and should be determined by the user.

Restrictions:

For Research Use only


Handling

Buffer:	PBS containing 0.02 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	Store the antibody undiluted at 2-8 °C.	



Immunofluorescence

Image 1. Immunocytochemistry of IKAP in A-20 cells with AP30402PU-N IKAP antibody at 1 μ g/ml.



Western Blotting

Image 2. Western blot analysis of IKAP in A-20 cell lysate with AP30402PU-N IKAP antibody at in (A) 0.5, and (B) 1 μ g/ml.