

Datasheet for ABIN4999763

anti-CDK11 antibody (AA 21-120) (AbBy Fluor® 680)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CDK11 (CDK11B)
Binding Specificity:	AA 21-120
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CDK11 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CDK11
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CDK11 (CDK11B)
Alternative Name:	CDK11 (CDK11B Products)
Background:	Synonyms: Galactosyltransferase associated protein kinase p58/GTA, CD11B_HUMAN,

Target Details

CDK11B, CDC2L1, PITSLREA, PK58, CDK11-p110, CDK11-p46, CDK11-p58, CDK11, CDK11 p110, CDK11 p46, CDK11 p58, CLK-1, p58, p58CDC2L1, p58CLK-1, p58 CLK-1, PK58, Cyclin-dependent kinase 11B, Cell division cycle 2-like protein kinase 1, Cell division protein kinase 11B, Galactosyltransferase-associated protein kinase p58/GTA, PITSLRE serine/threonine-protein kinase CDC2L1.

Background: Summary: This gene encodes a member of the p34Cdc2 protein kinase family. p34Cdc2 kinase family members are known to be essential for eukaryotic cell cycle control. This gene is in close proximity to CDC2L2, a nearly identical gene in the same chromosomal region. The gene loci including this gene, CDC2L2, as well as metalloprotease MMP21/22, consist of two identical, tandemly linked genomic regions which are thought to be a part of the larger region that has been duplicated. This gene and CDC2L2 were shown to be deleted or altered frequently in neuroblastoma with amplified MYCN genes. The protein kinase encoded by this gene could be cleaved by caspases and was demonstrated to play roles in cell apoptosis. Several alternatively spliced variants of this gene have been reported. [provided by RefSeq, Jul 2008].

Gene ID: 984

Pathways: [Regulation of Lipid Metabolism by PPARalpha, M Phase](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months