# antibodies - online.com







## anti-CINP antibody (Alexa Fluor 680)



Ovarvian

Overview	
Quantity:	100 μL
Target:	CINP
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CINP antibody is conjugated to Alexa Fluor 680
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human CINP

Immunogen:	KLH conjugated synthetic peptide derived from human CINP
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

l arget Details	
Target:	CINP
Alternative Name:	CINP (CINP Products)
Background:	Synonyms: CINP protein, CDK2-interacting protein, CINP, CINP_HUMAN, Cyclin dependent kinase 2 interacting protein, Cyclin-dependent kinase 2-interacting protein, MGC849.
	Background: Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin
	dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key

### **Target Details**

substrates involved in each phase of cell cycle progression. Specifically, Cdk2 interacts with Cyclins A, B1, B3, D, or E to control cell cycle progression. The Cyclin-dependent kinase 2-interacting protein (CINP) interacts with components of the replication complex and Cdk2 and Cdc7, thereby providing a functional and physical link between Cdk2 and Cdc7 during firing of the origins of replication. However, CINP is phopshorylated by Cdc7, but not by Cdk2. CINP also interacts with ATR-interacting protein and regulates ATR-dependent signaling, resistance to replication stress and G2 checkpoint integrity.

Gene ID:

51550

1 μg/μL

## **Application Details**

Concentration:

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol

bullet.	Aqueous burrered solution containing 0.0 fw rbs ( ph 7.4) with 1 % bsA, 0.05 % Proclinson and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months