antibodies -online.com





anti-CLK2 antibody (AA 401-499) (AbBy Fluor® 750)



Go to Product page

()	11/0	K\ /	iew
	\cup	'I V/I	$I \cap VV$

Quantity:	100 μL
Target:	CLK2
Binding Specificity:	AA 401-499
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CLK2 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CLK2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CLK2
Alternative Name:	CLK2 (CLK2 Products)

Target Details

Background:	Synonyms: CDC like kinase 2, CLK 2, CLK kinase, Clk2 Scamp3, Dual specicity protein kinase	
	CLK2, hCLK 2, hCLK2, MGC61500, Tu52, CLK2_HUMAN.	
	Background: CDC-like kinase 2 (CLK2) belongs to a family of autophosphorylating kinases	
	termed CLK (CDC2/CDC28-like kinases), which have been shown to phosphorylate serine- and	
	arginine-rich (SR) proteins of the spliceosomal complex, and to influence alternative splicing in	
	overexpression systems. Recent findings demonstrated that the CLK kinases activate PTP-1B	
	family members, and this phosphatase may be an important cellular target for CLK action.	
	Mutations in the CLK2 proteins affect organismal features such as development, behavior,	
	reproduction, and aging as well as cellular features such as the cell cycle, apoptosis, the DNA	
	replication checkpoint, and telomere length.	
Gene ID:	1196	
Pathways:	Regulation of Carbohydrate Metabolic Process	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	