

Datasheet for ABIN5000898

anti-CYR61 antibody (AA 250-300) (AbBy Fluor® 680)



Go to Product page

_			
()	V/C	rv	٨/

Background:

Quantity:	100 μL
Target:	CYR61
Binding Specificity:	AA 250-300
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYR61 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB)
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human Cyr61
	KLH conjugated synthetic peptide derived from human Cyr61 IgG
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	IgG Human
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity:	IgG Human Mouse,Rat,Dog,Cow,Sheep,Horse
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification:	IgG Human Mouse,Rat,Dog,Cow,Sheep,Horse

Synonyms: CCN1, GIG1, IGFBP10, Protein CYR61, CCN family member 1, Cysteine-rich

angiogenic inducer 61, Insulin-like growth factor-binding protein 10, IBP-10, IGF-binding protein 10, IGFBP-10, Protein GIG1, CYR61

Background: Promotes cell proliferation, chemotaxis, angiogenesis and cell adhesion. Appears to play a role in wound healing by up-regulating, in skin fibroblasts, the expression of a number of genes involved in angiogenesis, inflammation and matrix remodeling including VEGA-A, VEGA-C, MMP1, MMP3, TIMP1, uPA, PAI-1 and integrins alpha-3 and alpha-5. CYR61-mediated gene regulation is dependent on heparin-binding. Down-regulates the expression of alpha-1 and alpha-2 subunits of collagen type-1. Promotes cell adhesion and adhesive signaling through integrin alpha-6/beta-1, cell migration through integrin alpha-v/beta-5 and cell proliferation through integrin alpha-v/beta-3.

Gene ID: 3491

UniProt: 000622

Pathways: Positive Regulation of Endopeptidase Activity, Growth Factor Binding

Application Details

Application Notes: IF(IHC-P) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	