# antibodies -online.com





# anti-DAK antibody (AA 101-200) (Biotin)



Go to Product page

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
Target:	DAK
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DAK antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DAK	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

## **Target Details**

Target:	DAK
Alternative Name:	DAK (DAK Products)

## **Target Details**

ranger z erame		
Background:	Synonyms: Triokinase/FMN cyclase, TKFC, Triokinase/FMN cyclase, Bifunctional ATP-	
	dependent dihydroxyacetone kinase/FAD-AMP lyase (cyclizing), ATP-dependent	
	dihydroxyacetone kinase, DHA kinase, Glycerone kinase, Triokinase, Triose kinase, FAD-AMP	
	lyase (cyclizing), FAD-AMP lyase (cyclic FMN forming), FMN cyclase	
	Background: Catalyzes both the phosphorylation of dihydroxyacetone and of glyceraldehyde,	
	and the splitting of ribonucleoside diphosphate-X compounds among which FAD is the best	
	substrate. Represses IFIH1-mediated cellular antiviral response.	
Gene ID:	26007	
UniProt:	Q3LXA3	
Application Details		
Application Notes:	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	
Expiry Date:	12 months	