antibodies -online.com





anti-DAZ4 antibody (AA 51-150) (Alexa Fluor 750)



| () | 1 / | \sim | KI / | 110 | Νę |
|-----|-----|----------|------|-----|-------|
| | 1// | \vdash | I \/ | 1 ← | ٠// ٢ |
| | | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | DAZ4 | |
| Binding Specificity: | AA 51-150 | |
| Reactivity: | Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This DAZ4 antibody is conjugated to Alexa Fluor 750 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human DAZ4 | |
|-----------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Mouse | |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Pig,Horse | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | DAZ4 |
|-------------------|----------------------|
| Alternative Name: | DAZ4 (DAZ4 Products) |

Target Details

Background:

Synonyms: DAZ4, Deleted in azoospermia 4, Deleted in azoospermia protein 4.

Background: Spermatogenesis is the process by which male spermatogonia develop into mature spermatozoa. DAZ (deleted in azoospermia) are RNA-binding proteins that play an essential role in spermatogenesis. DAZ proteins influence the first stages of spermatogenesis and the maintenance of germ cell populations. DAZ proteins (DAZ1, DAZ2, DAZ3, DAZ4 and DAZ5) are encoded by separate genes on chromosome Y, each of which contain an AZFc domain in their coding region. DAZ proteins localize to the nucleus of spermatogonia, but relocate to the cytoplasm during meiosis. DAZ proteins contain an RRM (RNA recognition motif) domain that may regulate mRNA translation by binding to the 3?UTR. Deletions in the genes encoding DAZ proteins may cause azoospermia or oligospermia which can lead to male infertility. DAZ4 (deleted in azoospermia 4), also known as pDP1680 or pDP1681, is a 579 amino acid testis specific protein that contains nine DAZ-like domains and two RNA recognition motifs (RRM). DAZ4 exists as two alternatively spliced isoforms.

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 | |
|--------------------|--|--|
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |