antibodies -online.com





anti-POLI antibody (AA 51-150) (Alexa Fluor 680)



Quantity:	100 μL
Target:	POLI
Binding Specificity:	AA 51-150
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This POLI antibody is conjugated to Alexa Fluor 680

Application: Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Overview

Immunogen:	KLH conjugated synthetic peptide derived from human DNA Polymerase iota	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Dog	
Purification:	Purified by Protein A.	

Target Details

Target:	POLI
Alternative Name:	DNA Polymerase iota (POLI Products)

Target Details

Background:	Synonyms: DNA polymerase iota, Eta 2, Eta2, POLI, POLI_HUMAN, Polymerase DNA directed		
	iota, RAD 30B, RAD30 homolog B, RAD30B, RAD30B.		
	Background: DNA polymerase activity is essential for replication, repair, recombination and		
	mutagenesis. DNA polymerases can often bypass DNA lesions that block DNA replication,		
	thereby allowing the replication of damaged DNA. One such DNA polymerase is the distributive		
	enzyme DNA Pol i, which is encoded by the POLI gene. POLI is located on human chromosome		
	18q21.2, a region often implicated in the etiology of many human cancers. At thymine		
	templates, DNA Pol i is highly error-prone when replicating undamaged DNA in that it favors the		
	misincorporation of guanine over the correct nucleotide, adenosine. DNA Pol i also promotes		
	the replication of damaged DNA by misincorporating deoxynucleotides opposite DNA lesions.		
	DNA Pol i acts sequentially with DNA Pol , which is essential for damage-induced mutagenesis,		
	to complete the DNA lesion bypass. Therefore, replication involving DNA Pol i is likely to be		
	highly mutagenic.		
Gene ID:	11201		
Pathways:	DNA Damage Repair		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		

	_	
-	II	1:
-	-222	nnc
-	Hand	11111

Expiry Date:

12 months