

Datasheet for ABIN5001438

anti-DOK2 antibody (pTyr351) (AbBy Fluor® 680)



Overview

Overview	
Quantity:	100 μL
Target:	DOK2
Binding Specificity:	pTyr351
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DOK2 antibody is conjugated to AbBy Fluor® 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic phosphopeptide derived from mouse p56Dok2 around the phosphorylation site of (Tyr351)
Isotype:	IgG
Specificity:	This phosphorylation site is found at Tyr345 in Human. Due to the highly conserved nature of this sequence with DOK1, this antibody may react with DOK1 when it is phosphorylated at Tyr362 in Human and Tyr361 in Mouse and Rat.
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	DOK2
Alternative Name:	DOK2 (DOK2 Products)
Background:	Synonyms: DokR, Frip, Docking protein 2, Dok-related protein, Dok-R, Downstream of tyrosine kinase 2, IL-four receptor-interacting protein, p56(dok-2), Dok2 Background: DOK proteins are enzymatically inert adaptor or scaffolding proteins. They provide a docking platform for the assembly of multimolecular signaling complexes. DOK2 may modulate the cellular proliferation induced by IL-4, as well as IL-2 and IL-3. May be involved in modulating Bcr-Abl signaling. Attenuates EGF-stimulated MAP kinase activation.
Gene ID:	13449
UniProt:	070469

Application Details

Application Notes:	IF(IHC-P) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months