



[Go to Product page](#)

Datasheet for ABIN5001463

anti-RNF19A antibody (AA 621-720) (Alexa Fluor 750)

Overview

Quantity:	100 µL
Target:	RNF19A
Binding Specificity:	AA 621-720
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RNF19A antibody is conjugated to Alexa Fluor 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DORFIN/RNF19
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse
Purification:	Purified by Protein A.

Target Details

Target:	RNF19A
Alternative Name:	DORFIN/RNF19 (RNF19A Products)
Background:	Synonyms: Dorfin, Double ring finger protein, Double ring-finger protein, E3 ubiquitin-protein

Target Details

ligase RNF19A, p38, p38 protein, Ring finger protein 19, RING finger protein 19A, Ring IBR ring domain containing protein Dorfin, RN19A_HUMAN, RNF19, RNF19A, Ubce7ip2, UIP117, XYbp, AA032313, DKFZP566B1346.

Background: Dorfin is a multi-pass membrane, RING-IBR type, E3 ubiquitin-protein ligase. It is widely expressed with highest levels found in heart and ubiquitous expression throughout the central nervous system. Dorfin functions by accepting ubiquitin in the form of a thioester from UBC7 and UBC8 and then transferring it to the targeted substrates. Dorfin is responsible for ubiquitylating synphilin-1, CaSR and mutant variants of SOD-1, a protein at fault for familial ALS (amyotrophic lateral sclerosis). Dorfin physically interacts with VCP (Valosin-containing protein) via its C-terminus. Together these two proteins are associated with the formation of ubiquitylated inclusions (UBIs) that characterize many neurodegenerative disorders, such as Parkinson's disease and ALS. This association with UBIs suggests that Dorfin plays an important role in the disease process.

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months