# antibodies -online.com





## anti-PRUNE antibody (AA 201-300) (Alexa Fluor 750)



Go to Product page

| ( ) | 1/0 | r\ /1 | 014 |   |
|-----|-----|-------|-----|---|
| ( ) | ve  | I V I | -v  | V |
|     |     |       |     |   |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | PRUNE  |  |
| Binding Specificity: | AA 201-300   |  |
| Reactivity:          | Mouse  |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This PRUNE antibody is conjugated to Alexa Fluor 750   |  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human DRES17 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Mouse  |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit                   |
| Purification:         | Purified by Protein A.                                     |

### **Target Details**

| Target:           | PRUNE                   |
|-------------------|-------------------------|
| Alternative Name: | DRES17 (PRUNE Products) |

## Target Details

| Background:         | Synonyms: DRES17, Drosophila-related expressed sequence 17, hprune, HTCD37, Prune              |  |
|---------------------|--|--|
|                     | homolog, PRUNE like protein, TcD37 homolog, PRUNE_HUMAN.                                       |  |
|                     | Background: PRUNE, the human homologue of the Drosophila gene, is located in 1q21.3, a         |  |
|                     | region highly amplified in human sarcomas, malignant tumours of mesenchymal origin. Human      |  |
|                     | prune (h-prune), a phosphoesterase DHH family appertaining protein, physically interacts with  |  |
|                     | nm23-H1, a metastasis suppressor gene. h-prune is involved in cellular motility and metastasis |  |
|                     | formation. Metastatic breast cancers were found to overexpress h-prune, h-prune was also       |  |
|                     | found to be highly expressed in colorectal and pancreatic cancers. Hence h-prune is considered |  |
|                     | useful as a marker for tumor aggressiveness.   |  |
| Gene ID:            | 58497  |  |
| Application Details |  |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |  |
|                     | IF(IHC-F) 1:50-200   |  |
|                     | IF(ICC) 1:50-200   |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Concentration:      | 1 μg/μL  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and    |  |
|                     | 50 % Glycerol.   |  |
| Preservative:       | ProClin  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be            |  |
|                     | handled by trained staff only.   |  |
| Storage:            | -20 °C   |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.              |  |
| Expiry Date:        | 12 months  |  |
|                     |  |  |