

Datasheet for ABIN5001604

anti-DKC1 antibody (AA 81-190) (AbBy Fluor® 750)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	DKC1
Binding Specificity:	AA 81-190
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DKC1 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Dyskerin
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	DKC1
Alternative Name:	Dyskerin (DKC1 Products)
Background:	Synonyms: CBF5, CBF5 homolog, Cbf5p homolog, DKC 1, DKC, Dkc1, DKC1_HUMAN, DKCX,

Target Details

Dyskeratosis congenita 1, Dyskeratosis congenita 1 dyskerin, Dyskerin, H/ACA ribonucleoprotein complex subunit 4, NAP 57, NAP57, NAP-57, NOLA 4, NOLA4, Nopp140 associated protein of 57 kDa, Nopp140-associated protein of 57 kDa, Nucleolar protein family A member 4, Nucleolar protein NAP57, snoRNP protein DKC1, XAP 101, XAP101.

Background: Dyskerin (NAP57) associates with the chaperone protein Nopp140 and forms a small ribonucleoprotein particle with GAR1 (NOLA1), NHP2 (NOLA2) and Nop10 for the isomerization of uridine to pseudouridine (1). GAR1, NHP2 and dyskerin localize to the dense fibrillar component of the nucleolus and in nuclear Cajal bodies (2). The dyskerin gene maps to chromosome Xq28 (3). Missense mutations in the dyskerin gene interfere with normal nuclear localization of dyskerin and cause Dyskeratosis congenita (DKC) (4). DKC is a rare, X-linked bone marrow disorder characterized by cutaneous hyperpigmentation, dystrophy of the nails, atrophy of the testicles and leukoplakia of the oral mucosa. The GAR1 gene maps to chromosome 4q25 (5,6). The NHP2 gene maps to chromosome 5q35.3 and encodes a 155-amino acid protein (2,7).

Gene ID:	1736
Pathways:	Telomere Maintenance

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months