

Datasheet for ABIN5002114

anti-ERK1/2 antibody (pThr183, pTyr185) (AbBy Fluor® 750)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	ERK1/2 (MAPK1/3)
Binding Specificity:	pThr183, pTyr185
Reactivity:	Human, Mouse, Rat, Chicken
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ERK1/2 antibody is conjugated to AbBy Fluor® 750
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Flow Cytometry (FACS), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human ERK2 around the phosphorylation site of Thr183/Tyr185
Isotype:	IgG
Cross-Reactivity:	Chicken, Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Horse,Rabbit,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	ERK1/2 (MAPK1/3)
---------	------------------

Target Details

Alternative Name:	ERK1+ERK2 (MAPK1/3 Products)
Background:	Synonyms: ERK, p38, p4, p41, ERK2, ERT1, ERK-2, MAPK2, PRKM1, PRKM2, P42MAPK, p41mapk, p42-MAPK, ERK1, ERT2, ERK-1, PRKM3, P44ERK1, P44MAPK, HS44KDAP, HUMKER1A, p44-ERK1, p44-MAPK, Mitogen-activated protein kinase 1, MAP kinase 1, MAPK 1, Extracellular signal-regulated kinase 2, MAP kinase isoform p42, Mitogen-activated protein kinase 2, MAP kinase 2, MAPK 2, MAPK1 Background: The protein encoded by this gene is a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. The activation of this kinase requires its phosphorylation by upstream kinases. Upon activation, this kinase translocates to the nucleus of the stimulated cells, where it phosphorylates nuclear targets. Two alternatively spliced transcript variants encoding the same protein, but differing in the UTRs, have been reported for this gene.

Gene ID: 55945595

UniProt: [P28482](#)

Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months