# antibodies .- online.com





# anti-FANCM antibody (AA 831-930) (Alexa Fluor 680)



Go to Product page

$\sim$			
	N/P	r\/	i⊢₩

Quantity:	100 μL	
Target:	FANCM	
Binding Specificity:	AA 831-930	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FANCM antibody is conjugated to Alexa Fluor 680	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FANCM	
Isotype:	IgG	
Predicted Reactivity:	Human	
Purification:	Purified by Protein A.	

# **Target Details**

Target:	FANCM	
Alternative Name:	FANCM (FANCM Products)	
Background:	Synonyms: FAAP250, Fanconi anemia group M protein, Protein Hef ortholog,	

Background: Fanconi anemia (FA) is an autosomal recessive disorder characterized by bone marrow failure, birth defects and chromosomal instability. At the cellular level, FA is characterized by spontaneous chromosomal breakage and a unique hypersensitivity to DNA cross-linking agents. The thirteen FA proteins that have been characterized are important for regulating chromosomal stability and genome surveillance. Eight of these proteins, namely FANCA, FANCB, FANCC, FANCE, FANCF, FANCG, FANCL and FANCM, comprise the FA core complex, which catalyzes a key reaction in DNA repair: the monoubiquitination of FANCD2. FANCM (Fanconi anemia, complementation group M) is a member of the DEAD-box helicase family of proteins and contains a DEAH helicase domain and a nuclease domain. Localizing to chromatin fractions, FANCM is phosphorylated in a cell cycle-dependent manner and is believed to function as an anchor, recruiting the FA core complex to chromatin. Mutations in the gene encoding FANCM can lead to Fanconi anemia.

Gene ID: 57697

Pathways: DNA Damage Repair

## **Application Details**

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

#### Handling

50 % Glycerol.  Preservative: ProClin			
Buffer:  Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 a 50 % Glycerol.  Preservative:  ProClin  Precaution of Use:  This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should handled by trained staff only.  Storage:  -20 °C	Format:	Liquid	
Freservative: ProClin  Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should handled by trained staff only.  Storage: -20 °C	Concentration:	1 μg/μL	
Precaution of Use:  This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should handled by trained staff only.  Storage:  -20 °C	Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
handled by trained staff only.  Storage: -20 °C	Preservative:	ProClin	
	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Storage:	-20 °C	
	Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

1	1 11	١٠
$\vdash$	land	∥n∩
	ıarıa	

Expiry Date:

12 months