

## Datasheet for ABIN5002590

## anti-FEZ1 antibody (AbBy Fluor® 680)



Go to Product page

_						
	V	$\triangle$	r۱	/1	$\triangle$	Λ/
	' V '		ΙV			v v

Quantity:	100 μL		
Target:	FEZ1		
Reactivity:	Human, Rat, Mouse		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This FEZ1 antibody is conjugated to AbBy Fluor® 680		
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p))		
Product Details			
Immunogen:	KLH conjugated synthetic peptide derived from human FEZ1		
Isotype:	IgG		
Cross-Reactivity:	Human, Mouse, Rat		
Purification:	Purified by Protein A.		
Target Details			
Target:	FEZ1		
Alternative Name:	FEZ1 (FEZ1 Products)		
Background:	Synonyms: Fasciculation and elongation protein zeta 1, Fasciculation and elongation protein		
	zeta-1, Fez 1, Fez, Fez-like zinc finger protein, FEZ1, FEZ1_HUMAN, Fezl, Forebrain embryonic		
	zinc-finger like protein, Zpf312, Zygin 1, Zygin I, Zygin-1, Zygin1, ZyginI.		
	Background: Frequent loss of heterozygosity (LOH) at human chromosome 8p22-p21 is		

associated with various tumors including prostate and breast cancer. The 8p22-p21 region contains the FEZ1 gene, which is altered in tumors of the esophagus, prostate and breast. The FEZ1 protein (also known as leucine zipper putative tumor suppressor or LZTS1) contains a DNA-binding leucine zipper motif. FEZ1 is expressed in normal breast and prostate, but alterations in FEZ1 expression result in abnormal cell growth. The absence of FEZ1 expression is characteristic of breast and prostate cancer cell lines as well as primary breast and pro-state tumors. This absence of FEZ1 may be due to several factors, including mutations in the FEZ1 gene or hypermethylation of the CpG island flanking the FEZ1 promoter region. FEZ1 acts as a negative regulator of cell growth. During cell-cycle progression, FEZ1 localizes to microtubule components and is hyperphosphorylated by cAMP-dependent kinase.

Gene ID:

9638

IF(IHC-P) 1:50-200

## **Application Details**

Application Notes:

Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	