# antibodies -online.com





# anti-FGF6 antibody (AA 110-208) (Alexa Fluor 680)



Go to Product page

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
Target:	FGF6
Binding Specificity:	AA 110-208
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FGF6 antibody is conjugated to Alexa Fluor 680
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FGF6	
Isotype:	IgG	
Predicted Reactivity:	Predicted Reactivity: Human,Mouse,Rat,Dog,Cow,Horse,Rabbit	
Purification:	Purified by Protein A.	

# **Target Details**

Target:	FGF6
Alternative Name:	FGF6 (FGF6 Products)
Background:	Synonyms: FGF 6, FGF6, FGF-6, Fibroblast growth factor 6, Fibroblast growth factor 6

precursor	HBGF 6	HRGF6	HST	2 HST2
piccuisci	, , , , , , , , , , , , , , , , , , , ,		, , , , , ,	2,11012.

Background: FGF6 is a secreted heparin binding growth factor that is a member of the FGF family. Proteins of this family play a central role during prenatal development and postnatal growth and regeneration of a variety of tissues, by promoting cellular proliferation and differentiation. FGF6 is expressed in leukemia cell lines with platelet megakaryocytic differentiation potential. It signals through FGFR 1c, 2c, and 4. The FGF6 gene displays oncogenic transforming activity when transfected into mammalian cells. The mouse homolog of this gene exhibits a restricted expression profile predominantly in the myogenic lineage, which suggested a role in muscle regeneration or differentiation.

2251

Pathways: RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin

Signaling Pathway

## **Application Details**

Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	

IF(ICC) 1:50-200

Restrictions: For Research Use only

#### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	